





藥品異動(新藥)公告

108年01月03日
第108-01期
藥劑科發布

	舊藥樣品 舊藥袋商品名格式	新藥樣品 新藥袋商品名格式
藥品外觀		
藥品代碼	MTCVD	
藥品名稱 中文名 廠牌/健保碼	CARVEDILOL HEXAL TABLETS 卡菲蒂羅錠 台灣諾華大藥廠/ BC24967100	DILATREND TABLETS 25MG 達利全錠 羅氏大藥廠/BC20452100
劑型/規格	CARVEDILOL 25MG/tab	
外觀描述	白色錠劑，有「D5」、「BM」字樣，雙面皆有一字刻痕。	
治療分類	Anti-hypertension, for heart diseases, and other uses	
適應症	高血壓、鬱血性心臟衰竭。	
用法用量	本態性高血壓：開始治療前兩天的建議劑量為 12.5 公絲，一天一次。之後，建議劑量為 25 公絲，一天一次。必要時，接著以至少兩週之間隔，將劑量增加至每日最大建議劑量 50 公絲，一天一次或分次服用(一天兩次)。	
給藥途徑	口服	
常見副作用	暈眩、頭痛，通常是輕微的且尤其於治療的初期。	
給藥注意事項	1. 請按時測量血壓、心跳及定時回診 2. 除非醫師有特殊指示，否則不可擅自停藥。 3. 連續假期或旅行前請留意藥量是否足夠。 4. 有下列情形時，請事先告知醫師：曾對此藥過敏；懷孕、授乳期；有心臟、腎臟或肝臟疾病；嚴重的支氣管氣喘或慢性阻塞性肺病；或甲狀腺機能亢進。	



藥品異動(新藥)公告

懷孕/授乳分級	<p>Adverse events have been observed in animal reproduction studies. Adverse events, such as fetal/neonatal bradycardia, hypoglycemia, and reduced birth weight, have been observed following in utero exposure to beta-blockers as a class. Adequate facilities for monitoring infants at birth is generally recommended.</p> <p>/ It is not known if carvedilol is present in breast milk.</p> <p>According to the manufacturer, the decision to continue or discontinue breastfeeding during therapy should take into account the risk of infant exposure, the benefits of breastfeeding to the infant, and benefits of treatment to the mother. Breastfeeding is not recommended for women with heart failure related to peripartum cardiomyopathy due to the high metabolic demands of lactation and breastfeeding .</p>
備註	<p>羅氏大藥廠生產之 DILATREND TABLETS 25MG，供貨正常，故待台灣諾華大藥廠生產之 CARVEDILOL HEXAL TABLETS，舊品用罄，以新品替代使用</p>